

DIVERSITY OF BLADED BANGIALES (RHODOPHYTA) IN WESTERN
MEDITERRANEAN: RECOGNITION OF THE GENUS *THEMIS* AND
DESCRIPTIONS OF *T. BALLESTEROSII* SP. NOV., *T. IBERICA* SP. NOV. AND
PYROPIA PARVA SP. NOV.

Noemí Sánchez

Universitat de Girona, Facultat de Ciències, Departament de Ciències Ambientals,
17071 Girona, Spain

*Alba Vergés*¹

Universitat de Girona, Facultat de Ciències, Departament de Ciències Ambientals,
17071 Girona, Spain

César Peteiro

Instituto Español de Oceanografía (IEO), Centro Oceanográfico de Santander, 39080
Santander, Spain

Judith E. Sutherland

School of Biological Sciences, the University of Auckland, Private Bag 92019,
Auckland 1142, New Zealand

and Juliet Brodie

Natural History Museum, Department of Life Sciences, London SW7 5BD, UK

¹ Author for correspondence: email alba.verges@udg.edu

ABSTRACT:

The diversity of the bladed species of the red algal order Bangiales from the Iberian Mediterranean shores has been reassessed after a detailed study of this region. Prior to this study, 11 bladed species of Bangiales had been reported from Mediterranean waters: *Porphyra atropurpurea*, *P. cordata*, *P. coriacea*, *P. dioica*, *P. linearis*, *P. purpurea*, *P. umbilicalis*, *Pyropia leucosticta*, *Py. koreana* (as *Py. olivii*), *Py. elongata* (as *P. rosenburgii*) and *Py. suborbiculata*. A combined analysis of the nuclear nSSU and the plastid *rbcL* genes together with detailed morphological studies has confirmed the presence of species within the genera *Porphyra* and *Pyropia* and also revealed a third, undescribed genus, *Themis* gen. nov. *Porphyra linearis*, *Pyropia elongata* and the introduced *Pyropia koreana* had been previously listed for the Mediterranean and were recorded in the present study. An additional four species, including the introduced *Pyropia suborbiculata* and three new species: *Pyropia parva* sp. nov., *Themis ballesterosii* sp. nov. and *Themis iberica* sp. nov. were also observed. Hence, most of the *Porphyra* species traditionally reported along these shores were not reported in this survey. This new floristic Bangiales composition confirms the importance of the Mediterranean basin as a hotspot for biodiversity, possible endemics of ancient origin and high proportion of introductions. Our data also continue to confirm the extent of Bangiales diversity at regional and worldwide levels.

Key index words: Bangiales, Mediterranean, nSSU, *Porphyra*, *Pyropia*, *rbcL*, taxonomic revision, *Themis*